



EHD.ORG &gt; HEALTH

Loading...

## Facts on Drugs: Nicotine

Source: "Facts on Drugs: Nicotine." NIDA for Teens. National Institute on Drug Abuse. 19 Jan, 2005

[http://teens.drugabuse.gov/facts/facts\\_nicotine1.asp#top](http://teens.drugabuse.gov/facts/facts_nicotine1.asp#top).

Public Domain

### Table of Contents (TOC)

[Chapter 1: What Is It?](#)

[Chapter 2: What Are the Common Street Names?](#)

[Chapter 3: How Is It Used?](#)

[Chapter 4: How Many Teens Use It?](#)

[Chapter 5: What Are the Common Effects?](#)

[Chapter 6: Tobacco Kills](#)

[Chapter 7: Nervous or Depressed?](#)

[References](#)

TOC

Chapter 1: What Is It?

Back  
to Top

Nicotine is the drug in tobacco leaves. Whether someone smokes, chews, or sniffs tobacco, he or she is delivering nicotine to the brain. Each cigarette contains about 10 milligrams of nicotine. Nicotine is what keeps people smoking despite its harmful effects. Because the smoker inhales only some of the smoke from a cigarette and not all of each puff is absorbed in the lungs, a smoker gets about 1 to 2 milligrams of the drug from each cigarette [1]. A drop of pure nicotine would kill a person—in fact, nicotine can be used as a pesticide on crops. [2]

TOC

Chapter 2: What Are the Common Street Names?

Back  
to Top

You might hear cigarettes referred to as smokes, cigs, or butts. Smokeless tobacco is often called chew, dip, spit tobacco, or snuff.

Tobacco can be smoked in cigarettes, cigars, or pipes. It can be chewed or, if powdered, sniffed. An alternative to cigarettes is "bidis." Originally from India, bidis are hand-rolled. In the U.S., bidis are popular with teens because they come in colorful packages with flavor choices. Some teens think that bidis are less harmful than regular cigarettes. But bidis have even more nicotine, which may make people smoke more, causing them to be more harmful to the lungs than cigarettes. [3]

More than 3 and one-half million teens between the ages of 12 and 17 use tobacco—that's about 15 percent of teens that age. Of those, just over 3 million, or 13 percent, smoke cigarettes. In the U.S., 66.5 million people, or about 29 percent of the population, use tobacco. [9]

With each puff of a cigarette, a smoker pulls nicotine into his or her lungs where it is absorbed into the blood. In eight seconds, nicotine is in the brain, changing the way the brain works. This process happens so fast because nicotine is shaped like the natural brain chemical acetylcholine. Acetylcholine is one of many chemicals called neurotransmitters that carry messages between brain cells. Neurons have special spaces called receptors, into which specific neurotransmitters can fit, like a key fitting into a lock. Nicotine locks into acetylcholine receptors in different parts of the brain, rapidly causing changes in the body and brain. Nicotine raises the heart rate and respiration (breathing) rate, and causes more glucose, or blood sugar, to be released into the blood. This might be why smokers feel more alert after smoking a cigarette. [4],[6]

Nicotine also attaches to neurons (brain cells) that release a neurotransmitter called dopamine. Nicotine stimulates neurons to release unusually large amounts of dopamine. Dopamine stimulates the brain's pleasure and reward circuit, a group of brain structures called the limbic system involved in appetite, learning, memory, and feelings of pleasure. Normally, pleasurable feelings come from food, comfort, and the company of people you love. But smoking cigarettes causes a flood of dopamine in the smoker's brain. Its this flood of dopamine that gives the smoker intense feelings of pleasure. [6],[7]

Normally, neurons reabsorb neurotransmitters after they've done their job of signaling other brain cells. But cigarette smoke causes dopamine to stay in the spaces between neurons called synapses. Researchers don't yet know exactly what component of tobacco smoke blocks the reabsorption of dopamine into neurons. [7]

In 40 minutes, half the effects of nicotine are gone. So smokers get the urge to light up for another dose of the drug. After repeated doses of nicotine, the brain changes. To adjust to too much dopamine, the brain cuts production of the neurotransmitter and reduces the number of some receptors. Now, the smoker needs nicotine just to create normal levels of dopamine in his or her brain. Without nicotine, the smoker feels irritable and depressed. The smoker has trained the limbic system to crave tobacco. Think about how you long for a cold drink on a hot day. Or how you want a sandwich when you are hungry. Craving for tobacco is much stronger. [6]

These changes in the brain and body make nicotine highly addictive. Other addictive drugs of abuse, including heroin and cocaine, cause the same changes in the brain. [4]

While inhaling a cigarette, smokers are pulling more than nicotine into their lungs. Tobacco smoke contains more than 4,000 chemicals. Besides nicotine, the most dangerous chemicals in cigarette smoke are tar and carbon monoxide. Tar causes lung cancer, emphysema, and bronchial diseases. Carbon monoxide causes heart problems; smokers are at high risk for heart disease. [2],[6]

Smokers also have a dulled sense of smell and taste, reduced stamina for exercise and sports, and they smell of smoke. After smoking for a long time, smokers find that their skin ages faster and their teeth turn brown or discolored. [2]

TOC

## Chapter 6: Tobacco Kills

Back  
to Top

Each year, nearly half a million Americans die from tobacco use. [6] One of every six deaths in the United States is a result of smoking tobacco, making tobacco more lethal than all other addictive drugs combined. [2]

TOC

## Chapter 7: Nervous or Depressed?

Back  
to Top

Scientists are learning how tobacco and nicotine affect teen smokers. Studies going on for 25 years show a link between heavy teen smoking and fear of going outside (agoraphobia). Teens who smoke were 6 times more likely to get agoraphobia. And, teen smokers were 15 times more likely to have panic attacks than teens who did not smoke. Scientists think the reason is that nicotine hurts blood vessels to the brain, and also blocks air from the lungs. Whatever the reason, teen smokers are more likely to have panic attacks, anxiety disorders, and depression. [8]

TOC

## References

Back  
to Top

1. National Institute on Drug Abuse. NIDA Research Report—Nicotine Addiction (<http://www.drugabuse.gov/researchreports/nicotine/nicotine.html>): NIH Pub. No. 01-4342. Bethesda, MD: NIDA, NIH, DHHS. Printed July, 1998. Reprinted Aug., 1998.
2. National Institute on Drug Abuse. Mind Over Matter: The Brain's Response to Nicotine ([http://teens.drugabuse.gov/mom/mom\\_nic1.asp](http://teens.drugabuse.gov/mom/mom_nic1.asp)): NIH Pub. No. 00-3858. Bethesda, MD: NIDA, NIH, DHHS. Printed 1997. Reprinted 1998, 2000.
3. National Institute on Drug Abuse. NIDA Notes, vol 15, no 1: Teen Alternative to Cigarettes Has Higher Concentrations of Nicotine ([http://www.drugabuse.gov/NIDA\\_Notes/NNVol15N1/BBoard.html](http://www.drugabuse.gov/NIDA_Notes/NNVol15N1/BBoard.html)): Bethesda, MD: NIDA, NIH, DHHS. Retrieved June 2003.
4. National Institute on Drug Abuse. NIDA InfoFacts: Cigarettes and Other Nicotine Products

(<http://www.drugabuse.gov/Infobox/tobacco.html>): Bethesda, MD: NIDA, NIH, DHHS. Retrieved June 2003.

5. National Institute on Drug Abuse. NIDA Published Article: Parents: Nicotine is a Real Threat to Your Kids ([http://www.drugabuse.gov/Published\\_Articles/Nicotinethreat.html](http://www.drugabuse.gov/Published_Articles/Nicotinethreat.html)) Bethesda, MD: NIDA, NIH, DHHS. Retrieved June 2003.

6. National Institute on Drug Abuse. Brain Power! The NIDA Junior Scientists Program (<http://www.nida.nih.gov/JSP/JSP.html>): NIH Pub. No. 01-4575. Bethesda, MD: NIDA, NIH, DHHS. Printed 2001.

7. National Institute on Drug Abuse. Mind Over Matter: The Brain's Response to Drugs Teacher's Guide ([http://teens.drugabuse.gov/mom/tg\\_nic1.asp](http://teens.drugabuse.gov/mom/tg_nic1.asp)): NIH Pub. No. 020-3592. Bethesda, MD: NIDA, NIH, DHHS. Printed 1997. Reprinted 1998, 2002. Revised 2000.

8. National Institute on Drug Abuse. NIDA Notes, vol 16, no 1: Smoking May Lead to Anxiety Disorders in Adolescents and Young Adults ([http://www.drugabuse.gov/NIDA\\_Notes/NNVol16N1/Smoking.html](http://www.drugabuse.gov/NIDA_Notes/NNVol16N1/Smoking.html)): Bethesda, MD: NIDA, NIH, DHHS. Retrieved June 2003.

9. SAMHSA Office of Applied Studies. Results from the 2001 National Household Survey on Drug Abuse: Volume III; Detailed Tables Volume III-ABN; Prevalence Estimates and Sample Sizes (<http://www.samhsa.gov/oas/nhsda/2k1nhsda/vol1/chapter4.htm>): SAMHSA, DHHS. Retrieved June 2003. le 2.2A and Table 2.2B Source cited: SAMHSA Office of Applied Studies, National Household Survey on Drug Abuse 2000 and 2001

---

[Home](#) | [Terms of Use](#) | [Disclaimers](#) | [Privacy Policy](#) | [Copyright](#) | [Feedback](#) | [Endorsement Form](#) | [Site Map](#) | [Search](#)

[ehd@ehd.org](mailto:ehd@ehd.org)

*Inspiring health through education*

[Back to Top](#)

COPYRIGHT © 2001-2005, THE ENDOWMENT FOR HUMAN DEVELOPMENT, INC., ALL RIGHTS RESERVED. UNAUTHORIZED USE PROHIBITED.